

Diseases of concern in wild Australian marine turtles

Fact sheet

March 2023

Introductory statement

Of the seven marine turtle species in the world, six occur in Australian waters:

- flatback turtle (*Natator depressus*)
- green turtle (*Chelonia mydas*)
- hawksbill turtle (*Eretmochelys imbricata*)
- leatherback turtle (*Dermochelys coriacea*)
- loggerhead turtle (*Caretta caretta*)
- olive ridley turtle (*Lepidochelys olivacea*).

While nesting predominantly on the beaches of northern Australia, marine turtles can be found in waters around the entire continent. Females lay two to six clutches of eggs each year. The young hatch and make their way to the sea after an incubation period of approximately two months. After 20 to 50 years, depending on the species, the females return to the region they were born to continue the cycle ^[1].

Relatively little is known of the life of a wild marine turtle prior to reaching sexual maturity. This includes any potential disease impacts. Current disease knowledge is based on those individuals that wash up on shore, are inadvertently caught in fishing nets or otherwise enter care because of their debilitated state. The types of diseases affecting turtle populations, the prevalence in different species and populations and the effects these diseases might have on those populations remain largely unknown.

Aetiology and epidemiology

Significant diseases that have been reported in wild Australian marine turtles include:

- coccidiosis (see also WHA fact sheet "Disseminated coccidiosis in green turtles")
- fibropapillomatosis
- spirorchidiasis
- "soft shell syndrome" of green turtles.

Other diseases have been recognised in marine turtles outside Australia. These include **chlamydiosis** ^[2], **fungal** infections caused by a range of species including *Fusarium* sp. ^[3], *Aspergillus* sp. and *Beauveria* sp. ^[4], ***Cryptosporidium* sp.** infection ^[5], **herpesvirus** infections and **mycoplasmosis** ^[4].

Physical factors may also present major threats to sea turtle population viability. Non-infectious disease threats include **traumatic injuries** due to **boat strikes**, **entanglement** (most often plastic and/or fishing line) and **foreign body ingestion**. It is estimated that between 4866 and 14,600 turtles were captured in 8690 ghost nets sampled across northern Australia from 2005 to 2012 ^[1]. Globally, evidence shows at least 52% of all sea turtles have ingested plastic debris. A study found a 22% probability of mortality after ingesting one piece of plastic, which increased to 50% if 14 pieces had been ingested ^[6]. A range of microplastics were found in every one of over one hundred individuals from a global study of all seven of marine turtles species, including those from in Qld waters ^[7].

Ingestion can occur at all stages of a sea turtle's lifecycle but appears to be most frequent in juvenile and pelagic stages. Plastic, such as shopping bags, that physically resembles turtles' natural food (jellyfish) is ingested at a higher rate than other types ^[6, 8].

Coccidiosis in wild green turtles, caused by *Caryospora chelonae*, was first described in 1991, from an epidemic affecting large subadult and young adults in south-east Qld and northern NSW ^[9, 10]. Subsequent epizootics and sporadic cases have been recorded in Qld and NSW ^[9, 11]. The life cycle is unknown. Most cases are seen between September and February. In 2014, over 150 green turtles died of coccidiosis over a three-month period, from October to December. The majority of affected turtles were older animals, which is consistent with previous coccidiosis outbreaks ^[10-12]. See the WHA fact sheet "Disseminated coccidiosis in green turtles" for more information.

Fibropapillomatosis is strongly associated with chelonid herpesvirus 5 (ChHV-5), which is believed to be the aetiological agent. The disease occurs most commonly in the green turtle but has been recognised in all seven species of marine turtles ^[13]. Fibropapillomatosis has a worldwide distribution. Most cases occur in juveniles, but the disease has also been reported in sub-adults and adults. Spread is believed to be by horizontal transmission either through direct contact, exposure to virus-laden water ^[14] or by mechanical vectors such as marine leeches (*Ozobranchus* spp.) ^[15] or cleaner fish ^[16], that have been shown to carry the virus.

Prevalence of fibropapillomatosis in populations can vary from 1% to 92% with a greater prevalence seen in stranded turtles than in nesting populations. A study of green turtles in Qld found a higher prevalence in inshore seagrass habitats than on coral reefs, possibly because these areas contain a greater proportion of juvenile turtles ^[17]. In Moreton Bay in 1990 8% of 249 turtles surveyed had fibropapillomas. At Shoalwater Bay prevalence was 2 to 3% from 1988 to 1990. At Repulse Bay prevalence increased from 0% in 1988 to 22% in 1990. At the coral reef sites, no cases were seen in 1697 turtles sampled between 1988 and 1990. In 2011, investigation of a green turtle mortality event near Gladstone, Qld found 8% of 63 turtles with fibropapillomas ^[18].

Outside Australia, a study of 4407 green turtles and 401 hawksbill turtles in Indonesia identified a prevalence of 21.5% in green turtles and 0% in hawksbill turtles. Prevalence increased up to a curved carapace length of 85 cm and then decreased again, indicating a higher susceptibility in younger turtles ^[19]. Recent reports indicate a prevalence in Florida turtles of 20 to 50% and Hawaiian turtles of 60% ^[4].

Spirorchidiasis is caused by a range of digenetic trematodes or flukes from the genera *Spirorchis*, *Henotosoma*, *Unicaecum*, *Vasotrema*, *Hapalorhynchus*, *Learedius*, *Hapalotrema*, *Neospororchis*, *Amphiorcus*, *Carettacola* and *Haemoxenicon*. They are found throughout the world and have been recognised in green turtles, loggerhead turtles and hawksbill turtles ^[20].

The 1-3 mm adult spirorchids occur in the heart and great vessels. Eggs are laid, which may become trapped in terminal blood vessels producing a granulomatous response. Others may penetrate the gut and are passed via the faeces into the water. Once in the water they hatch to produce miracidia which penetrate the intermediate host, likely a mollusc. The miracidia develop into cercariae, which either leave the intermediate host or are eaten with it, and penetrate the skin or mucous membranes of the definitive chelonian host where they mature in the blood vessels ^[21].

For spirorchids, one study found a prevalence of 4.8% in 104 green turtles from turtle farms in Torres Strait, 33.3% in 15 turtles from an oceanarium on Magnetic Island, and 72.2% in 22 turtles found on coral reefs ranging from Torres Strait to Townsville ^[22]. A later study found a prevalence of 40.9% in 22 wild Qld turtles ^[23]. A more recent report identified spirorchids as a cause of mortality in 41.8% of 100 green turtles necropsied from southern Qld ^[24]. Seven of 12 green turtles necropsied during the Gladstone mortality event mentioned previously were positive for spirorchidiasis ^[18].

"Soft shell syndrome" refers to a syndrome reported in green turtles in Qld defined by carapace softness and ulceration. Cases were first reported in 2021 in the Great Sandy Marine Park on the Fraser Coast, following 2021-22 wet season floods that resulted in substantial environmental and pollution impacts to green turtle habitat. Cases continue to be reported into 2023, with the majority of clinical cases originating from the Great Sandy Marine Park, and a small number of cases from the Moreton Bay region ^[25].

One Health implications

Wildlife and the environment: more work is required to better understand the impacts of the diseases described in this fact sheet on marine turtle populations in Australia. Relatively little is known about the impact of both infectious and non-infectious disease on marine turtles. These diseases will likely add to the challenges on marine turtle populations from human-related factors such as harvesting, fishing and plastic pollution. None of the infectious diseases mentioned in this sheet are known to impact other wildlife species, however as several diseases have been linked to poor environmental conditions, it is likely that there may be flow on impacts to other wildlife species, and the environment in general.

Domestic animals: no direct impacts on domestic animals are recognised.

Humans: none of these diseases are recognised as zoonotic.

Clinical signs, pathology, diagnostic specimens and procedures

Trauma can cause soft tissue, musculoskeletal and shell damage, which can lead to secondary infections and air leakage into the coelom, resulting in buoyancy problems. Entanglement frequently results in drowning. Ingestion of plastic and other debris such as fishhooks can have a

range of effects. The items may just pass through the gastrointestinal tract, or they may cause gut impaction, buoyancy difficulties or perforation, potentially causing death. New techniques are being developed to help quantify microplastic in sea turtles [26, 27].

Trauma and external entanglements are evident on examination. Foreign body ingestion can be difficult to diagnose and relies on a combination of clinical signs and diagnostic imaging to identify fish hooks, impactions or gas in the coelom [8].

Coccidial infections can be asymptomatic. Clinical signs, when they occur, include diarrhoea, neurological signs, weakness, buoyancy difficulties and emaciation. Pathology centres mostly on a necrotising enteritis. Schizonts, with associated inflammation, can also be found in extra-intestinal sites, especially the brain. Diagnosis is by histopathology and faecal examination for coccidia.

Fibropapillomatosis-affected turtles are often in poor body condition. Cutaneous masses can be found anywhere on the skin, but most commonly appear on the front flippers, neck and periocular tissues. Depending on size and position the masses may interfere with vision and locomotion [4, 20].

Affected turtles may have a nonregenerative anaemia, lymphopaenia, neutrophilia, monocytosis, hypoproteinaemia, hypoalbuminaemia, elevated aspartate aminotransferase, hypocalcaemia, hyponatraemia and hypochloraemia [13, 28].

Grossly tumours may be up to 20 cm in diameter and appear as rubbery, white or grey, verrucous or smooth, sessile or pedunculated fibrous masses. Tumours may also be found internally, most commonly in the lungs, heart and kidney [4, 20].

Histologically they are classified as fibromas or myxofibromas. Skin tumours are characterised by marked hyperplasia of the epidermis (up to 30 cells thick instead of the normal four to seven cells) with vacuolation of the cytoplasm and ballooning degeneration of epithelial cells with rete ridges extending into the dermis. The dermis often consists of proliferating fibroblasts and mixed perivascular inflammatory cells. Occasional eosinophilic intranuclear inclusion bodies can be seen in epithelial cells. Visceral tumours are characterised by extensive fibroblast proliferation with few mitotic figures [20, 29].

Diagnosis is by observing the characteristic fibropapillomas grossly and by histologic confirmation. Techniques for cell culture and PCR of ChHV5 are being developed [30].

Spirochid infections are often asymptomatic. Clinical signs, when they occur, are non-specific and include anorexia, debilitation, emaciation, weakness, anaemia, inability to dive and neurological signs [20, 22].

Grossly, the presence of adult trematodes is associated with thrombosis and thickening and hardening of blood vessel walls [20, 22].

Histologically affected turtles demonstrate widespread granulomatous vasculitis, thrombosis and micro abscesses associated with extensive aggregations of spirochid eggs within vessels of the myocardium, aorta, lungs, liver, kidney, spleen, brain and gastrointestinal tract, with the spleen, lungs and gastrointestinal tract predominating. Granulomas consist of lymphocytes, macrophages, multinucleate giant cells, fibroblasts and occasional eosinophils [20, 22, 31, 32].

Diagnosis is by finding eggs via faecal examination or in tissues. Flotation techniques are not reliable and either a sedimentation technique or a direct smear is recommended for faecal examination. Adults can be difficult to find ^[21]. In one study, although eggs were detected microscopically in tissues, adults could not be found in 59.2% of 27 turtles examined ^[22]. An ELISA test was developed in the USA that can detect serum antibodies to the spirorchids *Learedius learedi*, *Haplotrema dorsopora* and *Carettacola hawaiiensis* in green turtles ^[33]. A PCR test is available that can be used to identify spirorchids in tissue samples ^[32].

Turtles with **soft shell syndrome** may range from severely emaciated to over-conditioned ^[34]. First reported cases involved large (>100kg) individuals, but smaller individuals (10-40kg) are now also involved ^[25]. There are areas of carapace softness and fluid accumulation under the scutes, and this progresses to loss of normal carapace pattern, sloughing of scutes, haemorrhage and ulceration of the dermis and exposure of carapace bone, that reflects a pathological diagnosis of severe coagulative necrosis. Secondary bacterial and fungal infections are common ^[34]. Lesions range from discrete focal areas to generalised carapace involvement and may also be seen on the head and flippers. Many cases also present with gastroenteritis and sepsis. There is currently no consistent evidence pointing to a primary pathogen ^[25], and there are no consistent biochemical or haematological changes, making prognostic evaluation and treatment challenging ^[34].

The cause of soft shell syndrome is not known. Hypothesised aetiologies include one or more of the following:

- nutritional disease due to flood-related change in composition, digestibility or nutritional quality of natural diet
- infectious disease due to a novel pathogen or increased impact of known pathogens
- environmental exposure to toxins
- altered turtle behaviours and vulnerability to entrapment and excessive solar exposure as a result of abnormal environmental conditions associated with flooding.

Differential diagnoses

The presence of diarrhoea is strongly suggestive of **ccidiosis** but could be seen with other causes of enteritis. The main differential diagnoses for neurological signs are **spirorchid** or **ccidial** infections. The gross appearance of **fibropapillomatosis** is characteristic. Clinical signs of **spirorchidiasis** are non-specific and could occur with a range of conditions including hypothermia, malnutrition or opportunistic bacterial, fungal and viral infections.

Prevention, control and treatment

There is no effective treatment for **ccidiosis**.

The only treatment for **fibropapillomatosis** is surgical removal of the masses. One study found no recurrence two months following removal ^[29]. Turtles with mild disease may recover spontaneously.

Praziquantel is reported to be effective at treating **spirorchid** infections ^[35, 36].

Individuals affected by **soft shell syndrome** may recover with supportive care and treatment of secondary infections, using a combination of topical and systemic medications ^[25, 34]. Topical

treatments include betadine baths (with soft debridement or complete excision of the affected scutes). Systemic antimicrobials (including antifungals) are administered based upon culture and sensitivity ^[34]. Treatment is more successful if animals arrive into care in good body condition ^[25].

Research

Despite considerable research being carried out on sea turtle biology and ecology relatively little is known about the impact of disease. The epidemiology of both coccidiosis and fibropapillomatosis is yet to be fully elucidated and will possibly involve multiple factors such as water quality, water temperature and the presence of biotoxins ^[13, 37]. The life cycles of both *C. cheloniae* and spirorchids also remain to be determined. The effects that these diseases may have on turtle populations are not known. Investigations into possible causes of **soft shell syndrome** are ongoing.

As well as infectious disease research, long term studies need to be carried out to determine the effects that physical factors such as climate change, boat strike, fishing line entanglement and plastic ingestion could have on sea turtle populations.

Surveillance and management

Wildlife disease surveillance in Australia is coordinated by the Wildlife Health Australia. The National Wildlife Health Information System (eWHIS) captures information from a variety of sources including Australian government agencies, zoo and wildlife parks, wildlife carers, universities and members of the public. Coordinators in each of Australia's States and Territories report monthly on significant wildlife cases identified in their jurisdictions. NOTE: access to information contained within the National Wildlife Health Information System dataset is by application. See the WHA website for more information <https://wildlifehealthaustralia.com.au/ProgramsProjects/eWHIS-WildlifeHealthInformationSystem.aspx>.

The finding of lesions consistent with **soft shell syndrome** is considered interesting and unusual and cases are logged in eWHIS. NSDI funding may be available to assist with investigation of these cases; please contact the WHA Wildlife Coordinator in your state or territory for further information.

There are over 150 marine turtle events recorded in eWHIS (25 March 2019), from all states and territories in Australia where marine turtles occur in the wild. Green turtles are the most common species represented, followed by hawksbills and loggerheads. The range of diagnoses is wide. Diagnoses recorded in at least five separate events include (in alphabetical order) cachexia (anorexia, emaciation, starvation), coccidiosis, obstruction, parasitism, pneumonia, septicaemia, trauma and trematodiasis. At least ten cases (primarily green turtles) also had significant evidence of foreign body ingestion and entanglement.

On behalf of the Qld Department of Environment and Science, SeaWorld is investigating marine animal strandings on the Qld coastline from Bundaberg to the NSW border. If you come across a freshly deceased marine turtle in this area, please contact Dr Claire Madden (Claire_madden@vrtp.com.au) for further information.

References and other information

1. Commonwealth of Australia (2017) Recovery Plan for Marine Turtles in Australia, D.o.t.E.a. Energy, Editor: Canberra. 154.
2. Homer BL, Jacobson ER, Schumacher J, and Scherba G (1994) Chlamydiosis in mariculture-reared green sea turtles (*Chelonia mydas*). *Veterinary Pathology*, **31**(1): 1-7.
3. Cabanes FJ, Alonso JM, Castella G, Alegre F, Domingo M, and Pont S (1997) Cutaneous hyalohyphomycosis caused by *Fusarium solani* in a loggerhead sea turtle (*Caretta caretta*). *Journal of Clinical Microbiology*, **35**(12): 3343-5.
4. St. Leger J (2019) Interesting conditions of sea turtle pathology. In 'Proceedings: Wildlife Health and Pathology Short Course.' (Ed J. Hall) pp. 118-123. (Australian Registry of Wildlife Health: Sydney).
5. Graczyk TK, Balazs GH, Work T, Aguirre AA, Ellis DM, Murakawa S, and Morris R (1997) *Cryptosporidium* sp. infections in green turtles, *Chelonia mydas*, as a potential source of marine waterborne oocysts in the Hawaiian Islands. *Applied and Environmental Microbiology*, **63**(7): 2925-7.
6. Wilcox C, Puckridge M, Schuyler QA, Townsend K, and Hardesty BD (2018) A quantitative analysis linking sea turtle mortality and plastic debris ingestion. *Scientific Reports*, **8**(1): 12536.
7. Duncan EM, Broderick AC, Fuller WJ, Galloway TS, Godfrey MH, Hamann M, . . . Omeyer LC (2019) Microplastic ingestion ubiquitous in marine turtles. *Global change biology*, **25**(2): 744-752.
8. Innis CJ (2019) Medical management and rehabilitation of sea turtles. In 'Mader's Reptile and Amphibian Medicine and Surgery.' (Eds S.J. Divers and S.J. Stahl) pp. 1382-1388. (Elsevier: St. Louis).
9. Gordon AN (2005) A necropsy-based study of green turtles (*Chelonia mydas*) in south-east Queensland. thesis School of Veterinary Science, University of Queensland
10. Gordon A, Kelly W, and Lester R (1993) Epizootic mortality of free-living green turtles, *Chelonia mydas*, due to coccidiosis. *Journal of Wildlife Diseases*, **29**(3): 490-494.
11. Chapman P, Owen H, Flint M, Traub RJ, Cribb TH, and Mills PC (2016) Molecular characterization of coccidia associated with an epizootic in green sea turtles (*Chelonia mydas*) in South East Queensland, Australia. *PLoS one*, **11**(2): e0149962.
12. Grillo T, Cox-Witton K, Gilchrist S, and Iglesias R (2015) Animal Health Surveillance Quarterly. Animal Health Australia: Canberra. 6-7.
13. Jones K, Ariel E, Burgess G, and Read M (2016) A review of fibropapillomatosis in green turtles (*Chelonia mydas*). *Veterinary Journal*, **212**: 48-57.
14. Work TM, Dagenais J, Balazs GH, Schettle N, and Ackermann M (2015) Dynamics of virus shedding and in situ confirmation of chelonid herpesvirus 5 in Hawaiian green turtles with fibropapillomatosis. *Veterinary Pathology*, **52**(6): 1195-201.
15. Greenblatt RJ, Work TM, Balazs GH, Sutton CA, Casey RN, and Casey JW (2004) The *Ozobranchus* leech is a candidate mechanical vector for the fibropapilloma-associated turtle herpesvirus found latently infecting skin tumors on Hawaiian green turtles (*Chelonia mydas*). *Virology*, **321**: 101-110.
16. Lu Y, Yu Q, Zamzow JP, Wang Y, Losey GS, Balazs GH, . . . Yanagihara R (2000) Detection of green turtle herpesviral sequence in saddleback wrasse *Thalassoma duperrey*: a possible mode of transmission of green turtle fibropapilloma. *Journal of Aquatic Animal Health*, **12**(1): 58-63.
17. Limpus CJ and Miller JC (1990) The occurrence of cutaneous fibropapillomas in marine turtles in Queensland, in Australian marine turtle conservation workshop, R. James, Editor. Queensland Department of Environment and Heritage and Australian Nature Conservation Agency. 186-188.
18. Flint M, Eden PA, Limpus CJ, Owen H, Gaus C, and Mills PC (2015) Clinical and pathological findings in green turtles (*Chelonia mydas*) from Gladstone, Queensland: investigations of a stranding epidemic. *Ecohealth*, **12**(2): 298-309.

19. Adnyana W, Ladds PW, and Blair D (1997) Observations of fibropapillomatosis in green turtles (*Chelonia mydas*) in Indonesia. *Australian Veterinary Journal*, **75**(10): 736-42.
20. Ladds P (2009) 'Pathology of Australian Native Wildlife.' (CSIRO Publishing: Melbourne).
21. Reavill DR, Schmidt RE, and Stevenson R (2004) Review of spirorchid flukes (Digenea: Spirorchidae) and three cases in freshwater turtles. In 'Proceedings Association of Reptilian and Amphibian Veterinarians'. Naples.
22. Glazebrook JS, Campbell RS, and Blair D (1989) Studies on cardiovascular fluke (Digenea: Spirorchidae) infections in sea turtles from the Great Barrier Reef, Queensland, Australia. *Journal of Comparative Pathology*, **101**(3): 231-50.
23. Glazebrook JS and Campbell RSF (1990) A survey of the diseases of marine turtles in northern Australia. II. Oceanarium-reared and wild turtles. *Diseases of Aquatic Organisms*, **9**: 97-104.
24. Flint M, Patterson-Kane JC, Limpus CJ, and Mills PC (2010) Health surveillance of stranded green turtles in southern Queensland, Australia (2006-2009): an epidemiological analysis of causes of disease and mortality. *Ecohealth*, **7**(1): 135-45.
25. Kay E. Soft shell syndrome in marine turtles in Qld. 2023.
26. Caron AG, Thomas CR, Berry KL, Motti CA, Ariel E, and Brodie JE (2018) Ingestion of microplastic debris by green sea turtles (*Chelonia mydas*) in the Great Barrier Reef: Validation of a sequential extraction protocol. *Marine pollution bulletin*, **127**: 743-751.
27. Caron AG, Thomas CR, Berry KL, Motti CA, Ariel E, and Brodie JE (2018) Validation of an optimised protocol for quantification of microplastics in heterogenous samples: A case study using green turtle chyme. *MethodsX*, **5**: 812-823.
28. Norton TM, Jacobson ER, and Sundberg JP (1990) Cutaneous fibropapillomas and renal myxofibroma in a green turtle, *Chelonia mydas*. *Journal of Wildlife Diseases*, **26**(2): 265-70.
29. Jacobson ER, Mansell JL, Sundberg JP, Hajjar L, Reichmann ME, Ehrhart LM, . . . Murru F (1989) Cutaneous fibropapillomas of green turtles (*Chelonia mydas*). *Journal of Comparative Pathology*, **101**(1): 39-52.
30. Mashkour N, Maclaine A, Burgess GW, and Ariel E (2018) Discovery of an Australian *Chelonia mydas* papillomavirus via green turtle primary cell culture and qPCR. *Journal of virological methods*, **258**: 13-23.
31. Gordon A, Kelly WR, and Cribb TH (1998) Lesions caused by cardiovascular flukes (Digenea: Spirorchidae) in stranded green turtles (*Chelonia mydas*). *Veterinary Pathology*, **35**(1): 21-30.
32. Chapman P, Owen H, Flint M, Soares Magalhaes RJ, Traub RJ, Cribb TH, . . . Mills PC (2017) Molecular epidemiology and pathology of spirorchiid infection in green sea turtles (*Chelonia mydas*). *International Journal for Parasitology: Parasites and Wildlife*, **6**(1): 39-47.
33. Graczyk TK, Aguirre AA, and Balazs GH (1995) Detection by ELISA of circulating anti-blood fluke (*Carettacola*, *Haplotrema*, and *Learedius*) immunoglobulins in Hawaiian green turtles (*Chelonia mydas*). *Journal of Parasitology*, **81**(3): 416-21.
34. Madden C. Soft shell turtle syndrome in turtles Qld. 2023.
35. Adnyana W, Ladds PW, and Blair D (1997) Efficacy of praziquantel in the treatment of green sea turtles with spontaneous infection of cardiovascular flukes. *Australian Veterinary Journal*, **75**(6): 405-7.
36. Jacobson ER, Harman GR, Maxwell LK, and Laille EJ (2003) Plasma concentrations of praziquantel after oral administration of single and multiple doses in loggerhead sea turtles (*Caretta caretta*). *American Journal of Veterinary Research*, **64**(3): 304-9.
37. Cardenas DM, Cucalon RV, Medina-Magues LG, Jones K, Aleman RA, Alfaro-Nunez A, and Cardenas WB (2019) Fibropapillomatosis in a green sea turtle (*Chelonia mydas*) from the southeastern Pacific. *Journal of Wildlife Diseases*, **55**(1): 169-173.

Acknowledgements

We are extremely grateful to the many people who had input into this fact sheet and would specifically like to thank Peter Holz, Claire Madden (SeaWorld), Emily Kay (Australia Zoo Wildlife Hospital) and Qld Dept of Environment and Science.

Updated: March 2023

To provide feedback on this fact sheet

We are interested in hearing from anyone with information on these conditions in Australia, including laboratory reports, historical datasets or survey results that could be added to the National Wildlife Health Information System. If you can help, please contact us at admin@wildlifehealthaustralia.com.au.

Wildlife Health Australia welcomes your feedback on fact sheets. Please email admin@wildlifehealthaustralia.com.au. We would also like to hear from you if you have a particular area of expertise and are interested in creating or updating a WHA fact sheet. A small amount of funding is available to facilitate this.

Disclaimer

This fact sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness, or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary or medical consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this fact sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at wildlifehealthaustralia.com.au

Email: admin@wildlifehealthaustralia.com.au

Or call +61 2 9960 6333